

## **D.I.R.E. Score: Patient Selection for Chronic Opioid Analgesia**

For each factor, rate the patient's score from 1-3 based on the explanations in the right hand column.

Score	Factor	Explanation
	<u>D</u> agnosis	1 = Benign chronic condition with minimal objective findings or no definite medical diagnosis. Examples: fibromyalgia, migraine headaches, nonspecific back pain. 2 = Slowly progressive condition concordant with moderate pain, or fixed condition with moderate objective findings. Examples: failed back surgery syndrome, back pain with moderate degenerative changes, neuropathic pain. 3 = Advanced condition concordant with severe pain with objective findings. Examples: severe ischemic vascular disease, advanced neuropathy, severe spinal stenosis.
	<u>I</u> ntractability	1 = Few therapies have been tried and the patient takes a passive role in his/her pain management process. 2 = Most customary treatments have been tried but the patient is not fully engaged in the pain management process, or barriers prevent (insurance, transportation, medical illness). 3 = Patient fully engaged in a spectrum of appropriate treatments but with inadequate response.
	<u>R</u> isk	(R = Total of P + C + R + S below)
	<u>P</u> ychological:	1 = Serious personality dysfunction or mental illness interfering with care. Example: personality disorder, severe affective disorder, significant personality issues. 2 = Personality or mental health interferes moderately. Example: depression or anxiety disorder. 3 = Good communication with clinic. No significant personality dysfunction or mental illness.
	<u>C</u> hemical Health:	1 = Active or very recent use of illicit drugs, excessive alcohol, or prescription drug abuse. 2 = Chemical copier (uses medications to cope with stress) or history of CD in remission. 3 = No CD history. Not drug-focused or chemically reliant.
	<u>R</u> eliability:	1 = History of numerous problems: medication misuse, missed appointments, rarely follows through. 2 = Occasional difficulties with compliance, but generally reliable. 3 = Highly reliable patient with meds, appointments & treatment.
	<u>S</u> ocial Support:	1 = Life in chaos. Little family support and few close relationships. Loss of most normal life roles. 2 = Reduction in some relationships and life roles. 3 = Supportive family/close relationships. Involved in work or school and no social isolation.
	<u>E</u> fficacy score	1 = Poor function or minimal pain relief despite moderate to high doses. 2 = Moderate benefit with function improved in a number of ways (or insufficient info – hasn't tried opioid yet or very low doses or too short of a trial). 3 = Good improvement in pain and function and quality of life with stable doses over time.

\_\_\_ Total score = D + I + R + E

Score 7-13: Not a suitable candidate for long-term opioid analgesia

Score 14-21: May be a candidate for long-term opioid analgesia

Source: Miles Belgrade, Fairview Pain & Palliative Care Center © 2005.



TMC

**DIRE Score: Patient Selection for Chronic Opioid Analgesia**

803198 R 6/27/2015

Patient Name: \_\_\_\_\_

DOB: \_\_\_\_\_  
Or label